

Isolated Vibrational Wavepackets in D_2^+ : Defining Superposition Conditions and Wavepacket Distinguishability

W. A. Bryan,^{1,*} J. McKenna,² E. M. L. English,¹ J. Wood,¹ C. R. Calvert,² R. Torres,³ D. S. Murphy,² I. C. E. Turcu,⁴ J. L. Collier,⁴ J. F. McCann,² I. D. Williams,^{2,†} and W. R. Newell^{1,‡}

¹*Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK*

²*Department of Pure and Applied Physics, Queen's University Belfast, Belfast BT7 1NN, UK*

³*Blackett Laboratory, Imperial College London, Prince Consort Road, London SW7 2BW, UK*

⁴*Central Laser Facility, Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, UK*

(Dated: February 2, 2008)

Tunnel ionization of room-temperature D_2 in an ultrashort (12 femtosecond) near infra-red (800 nm) pump laser pulse excites a vibrational wavepacket in the D_2^+ ions; a rotational wavepacket is also excited in residual D_2 molecules. Both wavepacket types are collapsed a variable time later by an ultrashort probe pulse. We isolate the vibrational wavepacket and quantify its evolution dynamics through theoretical comparison. Requirements for quantum computation (initial coherence and quantum state retrieval) are studied using this well-defined (small number of initial states at room temperature, initial wavepacket spatially localized) single-electron molecular prototype by temporally stretching the pump and probe pulses.

PACS numbers: 42.50.Hz, 82.53.Hn, 03.67.Lx

As with a broad spectrum of quantum systems, if a molecule laser-induced into a coherent superposition of vibrational states is free to evolve, a time-dependent wavepacket is created. The wavepacket spreads or de-phases significantly after the initial superposition, the result of the constraining potential surface anharmonicity. However, when the discrete eigenstates naturally rephase or revive, the original state of the system is almost fully recovered, and the initial quasi-classical periodicity returns. This phenomenon of wavepacket revival [1] has major implications in experimentally realisable time-evolving quantum systems. Through the precise optical preparation of a tailored wavepacket in a molecule, the wavepacket motion is imprinted on the dissociative states, observable with a probability governed by the amplitudes that contribute to the superposition.

In this Letter, we report the isolation and quantification of a vibrational wavepacket in a room-temperature ensemble of D_2^+ ions. We vary the initial coherence of the superposed states and distinguish structure in the wavepacket revival when the measurement condition is changed. An application of these results is as a single-internal degree-of-freedom *prototype* vibrational qubit (vibrational states, v , support the qubit), the building block of a molecular quantum computer (MQC). Quantum computation operations are not performed, rather we test the preparation and observation of a superposition. While MQC has been demonstrated in Li_2 [2], the D_2^+ system is considered as it is readily handled theoretically (numerical solution of the rotation-free 1D TDSE under the two-state approximation) [3] and generated experimentally (D_2 molecules at room temperature are predominantly in $v=0$, no state-selective cooling is required; short laser pulse defines a spatially localized initial state) [3, 4, 5].

Before implementing quantum computation (QC), five general ‘physical implementation’ requirements have been identified [6]: (1) the ability to initialize the qubit in a fiducial state, (2) the coherence time of the qubit is much longer than the operation times, (3) a universal set of logic operators are available, (4) a qubit-specific measurement capability exists and (5) the quantum system is well characterized and readily scalable. We quantify the fulfilment of requirements (1), (4) and (5) with a discussion of requirement (2).

Clearly requirement (3) lies at the heart of QC, which in future experiments will be attempted by optically controlling the phase [7] or amplitude [8] of each vibrational state, referred to as phase or amplitude manipulation. The additional degrees of freedom in a larger molecule could be employed [9], i.e. compound vibrational or bending modes, making scalable (multi-dimensional) QC feasible.

Tunnel ionization ($D_2 (v' = 0) + \text{pump} \rightarrow D_2^+ (v = 0, 1, 2, \dots) + e^-$) when the Keldysh parameter $\gamma < 1$ [10] creates a vibrational wavepacket on the $1s\sigma_g$ potential surface of D_2^+ [3, 4, 5], and is near-instantaneous so is treated as a ‘vertical’ transition. The configuration of the potential energy surfaces is such that ionization only transfers population around the inner turning point of the wavepacket motion [3, 4, 5]. In a phase manipulation MQC, such a transition fulfils requirement (1), as all v -states are initiated with essentially zero phase difference. In an amplitude manipulation MQC, requirement (1) is similarly fulfilled as all accessible v -states have a known (Franck-Condon) amplitude distribution [11]. However, in practice an ensemble of molecules may be used to generate meaningful signal, where it becomes important that the pulse length is sufficiently short to retain coherence in the ensemble.




FIG. 1: (Color online) Comparison between experiment and theory showing vibrational and rotational revivals. (a) D^+ PD (top) and DI (bottom) yield as a function of wavepacket evolution time, Δt , and ion kinetic energy, recorded at room temperature. (b) Integrated PD yield (thick black) contains vibrational and rotational revivals (top). The theoretical prediction (green / thick grey curve) shows remarkable agreement with the experimental results when rotational dynamics are excluded (thin black curve). (c) The calculated time-dependent probability distribution. (d) The calculated impulsive alignment of D_2 induced by the pump pulse in a room-temperature distribution of rotational levels. (e) The predicted Franck-Condon (FC) distribution. (f) Fourier Transform of the experimental PD yield. The frequency components present in the vibrational wavepacket are beats between vibrational states v and $v \pm 1$. The Fourier amplitudes thus return a measure of vibrational population.

Observing the vibrational wavepacket evolution on the $1\sigma_g$ ground potential surface is possible by ‘collapsing’ the wavepacket to an observable state with a second laser pulse. Photodissociation (PD) $D_2^+ + probe \rightarrow D^+ + D$ breaks the internuclear bond via photon coupling the $1\sigma_g$ ground and $2p\sigma_u$ first excited states [12]. A second route to observing wavepacket evolution is through double ionization (DI) $D_2^+ \rightarrow D^+ + D^+ + e^-$, requiring a higher laser intensity as a second ionization event must occur [12]. PD and DI are distinguishable by the kinetic energy released during bond fracture. Both PD and DI are enhanced when the wavepacket is at the outer turning point $\simeq 3.5$ a.u., [3, 4, 5], and the internuclear potential energy surfaces naturally act as a quantum ‘shutter’ or ‘barrier’ depending on whether they are laser-coupled or not.

The experimental set-up is an extension of one described elsewhere [3, 4, 13]: intense (4×10^{14} Wcm $^{-2}$) ultrashort (12 femtosecond, $1 \text{ fs} = 10^{-15} \text{ s}$) near infrared (800 nm) laser pulses are used to create (pump) and observe (probe) the vibrational wavepacket. A co-linear low-dispersion Mach-Zehnder interferometer capable of supporting the pulse bandwidth ($> 100 \text{ nm}$) generates the time-separated pump and probe, allowing a delay $0 \leq \Delta t \leq 160$ picoseconds between the pump and probe with a repeatable resolution of $\simeq 300$ attosecond ($1 \text{ as} = 10^{-18} \text{ s}$). An alternative method for generating two or three time-delayed pulses has been demonstrated [14] by transmitting a $\simeq 10 \text{ fs}$ pulse through a disc-double annulus optic allowing two variable delays of hundreds of fs if the two annuli are radially rotated. After focussing with a spherical mirror, the interaction volumes generated by our Mach-Zehnder configuration overlap perfectly, whereas in the disc-annulus configuration, a time-varying spatial overlap is produced. Furthermore, the dispersion of the Mach-Zehnder configuration does not change with delay. D^+ fragments from PD and DI, Fig. 1(a), are detected in an ion time-of-flight mass spectrometer containing a $250\mu\text{m}$ aperture, thus signal can only result from PD or DI when the molecular axis is parallel to the probe polarization and detector axis. The aperture also defines a quasi-isointensity source in the focal volume.

In addition to creating a D_2^+ vibrational wavepacket, few-cycle pulses impulsively align deuterium [13, 14], thus the revival of a rotational wavepacket is expected. We isolate the vibrational and rotational motions, which is only possible for a limited range of pump pulse durations as discussed later. As shown in Fig. 1(a), in the region $450 \leq \Delta t \leq 650 \text{ fs}$, both PD and DI yields exhibit a $\simeq 24 \text{ fs}$ revival modulation with maximum contrast at 550 fs, the result of vibrational revival, consistent with recent findings [3, 4, 5]. However, between $250 \leq \Delta t \leq 350 \text{ fs}$ a rotational revival is also observed, comparable to recent observations [13, 14] which will also underlie the vibrational revival around 550 fs. Both the rotational and vibrational revivals are evident in the integrated PD yield as a function of Δt , Fig. 1(b) (top).

A major finding of the present work concerns the separation of these two wavepackets. The isolation of the vibrational wavepacket requires confirmation that two distinct populations in D_2 and D_2^+ exist following the pump. In recent work by the authors [13] (see also [14]), we quantify the rotational wavepacket in D_2 by Fourier transformation of the high-energy time-dependent DI signal recorded under identical experimental conditions. We observe peaks (with an accuracy of 5%) in the frequency spectrum at 5.5 (0), 9.2 (1), 12.8 (2), 16.4 (3), 20.1 (4) and 23.7 (5) THz, where parenthesis indicate the $J : J + 2$ rotational state beat in the D_2 molecule. There is no evidence for a rotational wavepacket in D_2^+ , which would contribute peaks in the frequency spectrum at 2.7 (0),




FIG. 2: (Color online) Controlling the initial superposition in an isolated vibrational wavepacket. (a) Experimental PD yield as the pump duration is increased (thin black curve). The rotational wavepacket has been removed from the dataset as discussed. The shape of the revival indicates the superposition degradation as the pump is stretched, illustrated by comparison with a theoretical prediction (thick magenta / grey curves). (b) The predicted vibrational wavepacket after averaging over the changing Gaussian pump duration.

4.5 (1), 6.3 (2), 8.1 (3), 9.9 (4), 11.7 (5), 13.5 (6), 15.3 (7) and 17.1 (8) THz. This is confirmed in the present work. Furthermore, the vibrational wavepacket can only exist in the D_2^+ molecular ion generated in the pump, as the pump duration is greater than the quasi-classical period (11 fs) of D_2 as recently investigated [15].

Thus we find that the pump either creates a vibrational wavepacket in D_2^+ (rotating in a non-time-correlated manner with respect to the pump with a room-temperature distribution of J-states) or a rotational wavepacket in D_2 (vibrating incoherently in an impulsively modified distribution of J-states). Solving the TDSE for the vibrational [3] (Fig. 1(c)), and rotational [13] (Fig. 1(d)) degrees of freedom, we quantify these motions. We take the integrated PD yield as a function of Δt (Fig. 1(b) top), and subtract the probability of finding the rotational wavepacket within the detector acceptance, Fig 1(d). The integrated PD yield (arb. units) is scaled relative to the rotational wavepacket probability until the signature of the rotational wavepacket ($250 \leq \Delta t \leq 350$ fs) is totally suppressed. The resulting PD yield contains the pure vibrational wavepacket, as shown in Fig. 1(b) (bottom). The PD yield is modelled by numerically propagating the initially localized state (FC principle) then calculating the time dependent coupling of the $1s\sigma_g$ and $2p\sigma_u$ states [3]. The vibrational wavepacket in the molecular ion is clearly well-described by this relatively simple treatment, thus requirement (5) is partially fulfilled. If evidence for strong rotational-vibrational coupling had been found, this would not be the case.

Fourier transformation of the raw time-dependant PD yield, Fig 1(b), recovers the beat frequencies of the vibrational levels with the associated amplitudes that form

the vibrational wavepacket, therefore we have a method for recovering the population of the vibrational states. As is apparent from Fig. 1(e) and (f), the measured distribution of vibrational beat frequencies agrees with the expected FC distribution of states for $v > 4$. For states $v < 4$, negligible beating is observed. This is a result of the probe pulse being of insufficient intensity to open the three-photon crossing [12], thus the vibrational wavepacket ‘below’ $v = 4$ remains bound. Our ‘quantum shutter’ is therefore selective, however the principle is sound, since given a probe intensity $> 6 \times 10^{14}$ Wcm $^{-2}$ all v states will be accessed. It should be noted that the states $0 < v < 4$ must be populated, otherwise the quasi-classical period of D_2^+ would not be observed: lack of population in these states would increase the vibrational revival periodicity. In a phase manipulation MQC, the relative phase can be temporally resolved for occupied v -states; while in an amplitude manipulation MQC, the relative amplitudes of the constituent v -states can also be recovered by Fourier transform analysis, Fig. 1(f). Returning to QC requirement (4), the ability to determine the final state in our measurement is met. Destruction of the prototype qubits is necessary to read the final state, however it is not necessary to destroy the wavepacket to manipulate. By utilizing nondestructive pulses, it should be possible to modify the wavepacket while maintaining the superposition.

To extend our understanding of fulfilling requirements (1) and (4), we quantify variations in the initial vibrational wavepacket coherence. We also study wavepacket distinguishability, the faculty to temporally discern subsequent oscillations of the vibrational wavepacket, particularly around revival. Both are macroscopically achieved through independently varying the pump and probe durations by changing the dispersion (hence chirp) in either arm of the interferometer.

Fig. 2(a) shows the integrated PD yield in the vicinity of the vibrational revival as the pump duration, τ_{pu} , is increased from 14 to 28 fs while the probe duration, τ_{pr} , is fixed at 13 fs. The vibrational wavepacket in D_2^+ was isolated from the rotational signature in D_2 as described earlier. The contrast of the vibrational revival structure decreases as the pump duration increases, the consequence of loss of coherence in the initial superposed state. The experimental revival structure (thin black curve, Fig. 2(a)) is compared to the theoretical wavepacket (Fig. 1(c)) averaged over the pump pulse, Fig. 2(b), and integrated $3 \leq R \leq 10$ a.u., then overlaid (thick magenta / grey curves) on the experimental data in Fig. 2(a). The resultant magenta curves show an excellent agreement with the decohering isolated vibrational wavepacket, clearly demonstrating the role of coherence in the initial pump pulse.

In Fig. 2(a), it is apparent that even for $\tau_{pu} = 28$ fs, some vibrational coherence remains when τ_{pu} is longer than the revival oscillatory period, 24fs. This residual




FIG. 3: (Color online) Distinguishing the reviving vibrational wavepacket as the measurement condition is changed. (a) The sudden loss of revival fidelity when the probe duration exceeds 40 fs, the consequence of the temporal structure becoming indistinguishable to the probe. From top to bottom, the probe durations are indicated sequentially. Comparing a (b) 15 fs, (c) 30 fs and (d) 45 fs probe to the revival structure, the facility to discern subsequent oscillations (i.e. wavepacket distinguishability) of the D_2^+ vibrational revival is compromised.

structure is due to the highest vibrational states, $v > 9$ which have the longest periods and an absolute population of less than 4%. Consequently, care must be taken in defining the initial state of a vibrational qubit as states with an apparently negligible population clearly influence wavepacket evolution.

By increasing τ_{pr} from 14 to 60 fs while keeping the pump fixed at $\tau_{pu} = 14$ fs (thus the vibrational wavepacket is identical to Fig. 2(b), bottom, throughout), the amplitude of the revival changes suddenly as τ_{pr} exceeds 40 fs. Since a 15 fs probe, Fig. 3(b), is short compared to the revival, oscillations are observed with high fidelity. At 30 fs, Fig. 3(c), a comparable fidelity is also achieved, as subsequent revivals are still distinguished. However, a 45 fs probe, Fig. 3(d), cannot distinguish any two revival maxima, hence the oscillatory structure is suddenly lost when τ_{pr} approaches double the quasi-classical period of D_2^+ . We can therefore define a wavepacket distinguishability condition, which is less strict than for creation, depending on the measurement regime. If a time-dependent measurement is performed at revival (as required in the phase manipulation), τ_{pr} must be shorter than double the quasi-classical period to ensure total wavepacket collapse. Alternatively, to guarantee that all the wavepacket is measured in an energy-resolved scheme (as required in the amplitude manipulation), τ_{pr} of at least double the period of the highest populated state is required. Furthermore, the intensity requirements discussed earlier must also be adhered to.

The pump-variation observations indicate a promising future for controlling the initial superposition con-

ditions. Following the optical preparation of an isolated vibrational wavepacket in a single molecule, a genetic algorithm will determine the intermediate time- and frequency-structured laser pulse [16] required to achieve a particular target state (either by phase or amplitude manipulation). This intermediate ‘operator’ pulse must be low intensity so as not to fracture the internuclear bonds, rather to guide wavepacket evolution. The high intensity ultrashort probe pulse will then fracture the internuclear bonds, allowing the result of the operation to be read.

To perform logic operations and apply successful error correction schemes requires a coherence time $\simeq 10^4$ times the interaction time, t_I , requirement (2). Here, ionization defines $t_I \simeq 1.3$ fs thus if the wavepacket survives to $\simeq 10$ ps with high fidelity, QC is possible. Our measurements have exceeded 2 ps, therefore this constraint does not appear technologically limiting. The MQC will be scaled by increasing the number of vibrational modes supporting the qubit, accomplished in a larger molecule. Communication between states will then be controlled optically by distorting the electronic potential surface through the Stark-shift.

In a broader application, a pump of a shorter duration than the vibrational motion establishes a time-dependent variation of the refractive index in a Raman-active molecular medium, as demonstrated in SF_6 [17]. By employing the vibrational mode of a hydrogenic molecular ion, it may be possible to spectrally broaden and self-compress, potentially to the single-cycle level.

The experiment was carried out at the ASTRA Laser Facility, STFC RAL, UK. The assistance of J. M. Smith, E. J. Divall, K. Ertel, O. Chekhlov, C. J. Hooker and S. Hawkes is gratefully acknowledged. This work was funded by the Engineering and Physical Sciences Research Council (UK). RT acknowledges the Spanish Department of State of Education and Universities, and the European Social Fund.

* Also at Central Laser Facility, Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, UK; Electronic address: w.bryan@ucl.ac.uk

† Electronic address: i.williams@qub.ac.uk

‡ Electronic address: w.r.newell@ucl.ac.uk

- [1] R. W. Robinet, Phys. Rep. **392** 1 (2004).
- [2] J. Vala *et al*, Phys. Rev. A **66** 062316 (2002).
- [3] D. S. Murphy *et al*, J. Phys. B **40**, S358 (2007).
- [4] J. McKenna *et al*, J. Phys. Conf. Ser. **58**, 375 (2007).
- [5] Th. Ergler *et al*, Phys. Rev. Lett. **97** 193001 (2006).
- [6] D. P. DiVincenzo, Fortschritte der Physik **48**, 771 (2000).
- [7] K. F. Lee, D. M. Villeneuve, P. B. Corkum and E. A. Shapiro, Phys. Rev. Lett. **93** 233601 (2004).
- [8] H. Niikura, D. M. Villeneuve and P. B. Corkum, Phys. Rev. Lett. **92** 133002 (2004).
- [9] C. M. Tesch and R. de Vivie-Riedle, Phys. Rev. Lett. **89**

- 157901 (2002).
- [10] V. S. Popov, Physics - Uspekhi **47** 855 (2004).
 - [11] X. Urbain *et al*, Phys. Rev. Lett. **92** 163004 (2004).
 - [12] J. H. Posthumus, Rep. Prog. Phys. **67**, 623 (2004).
 - [13] W. A. Bryan *et al*, Phys. Rev. A **76** 023414 (2007).
 - [14] K. F. Lee *et al*, J. Phys. B **39** 4081 (2006).
 - [15] Th. Ergler *et al*, Phys. Rev. Lett. **97** 103004 (2006).
 - [16] R. J. Levis, G. M. Menkir and H. Rabitz, Science **292** 709 (2001).
 - [17] N. Zhavoronkov and G. Korn, Phys. Rev. Lett. **88** 203901 (2002).